

Tel: (519) 489-7195, (800) 836-8089

Fax: (519) 231-0140, (877) 221-3515

Email: info@biomatik.com http://www.biomatik.com



IFI44L Antibody

CAC08158
Interferon-induced protein 44-like
Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Q53G44
Recombinant Human Interferon-induced protein 44-like protein
(206-307AA)
Rabbit
Human, Mouse
ELISA, WB, IF; Recommended dilution: WB:1:500-1:5000, IF:1:50-1:200
Exhibits a low antiviral activity against hepatitis C virus.
Liquid
Non-conjugated
Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
>95%, Protein G purified
IgG
Polyclonal
Interferon-induced protein 44-like, IFI44L, C1orf29
Human
Immunology
IFI44L




Biomatik

Tel: (519) 489-7195, (800) 836-

8089

Fax: (519) 231-0140, (877) 221-3515

Email: info@biomatik.com http://www.biomatik.com



Western Blot


Positive WB detected in: HepG2 whole cell lysate, A549 whole cell lysate, NIH/3T3 whole cell lysate

All lanes: IFI44L antibody at 2.5µg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 52, 19 kDa Observed band size: 52 kDa



Immunofluorescent analysis of HepG2 cells using CAC08158 at dilution of 1:100 and Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L)