

BiomatikTel: (519) 489-7195, (800) 836-8089
Fax: (519) 231-0140, (877) 221-3515
Email: info@ biomatik.com https://www.biomatik.com

anti- TRADD antibody

Product Information

Catalog No.: CAF50358

Size: 100µg Form: liquid

Purification: Immunogen affinity purified

≥95% as determined by SDS-PAGE Purity:

Host: Rabbit

Clonality: polyclonal

Clone ID: None IsoType: IgG

PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 Storage:

months (Avoid repeated freeze / thaw cycles.)

Background

The nuclear form acts as a tumor suppressor by preventing ubiquitination and degradation of isoform p19ARF/ARF of CDKN2A by TRIP12: acts by interacting with TRIP12, leading to disrupt interaction between TRIP12 and isoform p19ARF/ARF of CDKN2A(By similarity). Adapter molecule for TNFRSF1A/TNFR1 that specifically associates with the cytoplasmic domain of activated TNFRSF1A/TNFR1 mediating its interaction with FADD. Overexpression of TRADD leads to two major TNF-induced responses, apoptosis and activation of NF-kappa-B.

Immunogen information

TNFRSF1A-associated via death domain Immunogen:

Hs.89862, TRADD Synonyms:

34 kDa Observed MW: Uniprot ID: Q15628

Application

Reactivity: Human, Rat


Tested Application: ELISA, WB, IHC


Recommended dilution: WB: 1:200 - 1:2000; IHC: 1:50 - 1:500

Image:




Biomatik Tel: (519) 489-7195, (800) 836-8089 Fax: (519) 231-0140, (877) 221-3515 Email: info@ biomatik.com https://www.biomatik.com





Immunohistochemistry of paraffin-embedded rat liver tissue slide using CAF50358(TRADD antibody at dilution of 1:200



Jurkat cells were subjected to SDS PAGE followed by western blot with CAF50358(TRADD Antibody) at dilution of 1:1000