

BiomatikTel: (519) 489-7195, (800) 836-8089
Fax: (519) 231-0140, (877) 221-3515
Email: info@ biomatik.com https://www.biomatik.com

anti- TRAF6 antibody

Product Information

Catalog No.: CAF50359

Size: 100µg Form: liquid

Purification: Immunogen affinity purified

≥95% as determined by SDS-PAGE Purity:

Host: Rabbit

Clonality: polyclonal

IgG IsoType:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20° C for 24 Storage:

months (Avoid repeated freeze / thaw cycles.)

Background

E3 ubiquitin ligase that, together with UBE2N and UBE2V1, mediates the synthesis of 'Lys-63'linked-polyubiquitin chains conjugated to proteins, such as IKBKG, IRAK1, AKT1 and AKT2. Also mediates ubiquitination of free/unanchored polyubiquitin chain that leads to MAP3K7 activation. Leads to the activation of NF-kappa-B and JUN. May be essential for the formation of functional osteoclasts. Seems to also play a role in dendritic cells(DCs) maturation and/or activation. Represses c-Myb-mediated transactivation, in B-lymphocytes. Adapter protein that seems to play a role in signal transduction initiated via TNF receptor, IL-1 receptor and IL-17 receptor. Regulates osteoclast differentiation by mediating the activation of adapter protein complex 1(AP-1) and NF-kappa-B, in response to RANK-L stimulation. Together with MAP3K8, mediates CD40 signals that activate ERK in B-cells and macrophages, and thus may play a role in the regulation of immunoglobulin production.

Immunogen information

Immunogen: TNF receptor-associated factor 6

Synonyms: RING finger protein 85, RNF85, TRAF6

Observed MW: 60 kDa Uniprot ID: Q9Y4K3

Application

Reactivity: Human, Mouse, Rat; other species are not tested. Please decide the

specificity by homology


ELISA, WB, IHC, IF **Tested Application:**

Recommended dilution: WB: 1:500 - 1:2000; IHC: 1:50 - 1:200; IF: 1:50 - 1:200


Image:



Biomatik Tel: (519) 489-7195, (800) 836-8089 Fax: (519) 231-0140, (877) 221-3515 Email: info@ biomatik.com https://www.biomatik.com



Immunohistochemistry of paraffin-embedded human colon tissue slide using CAF50359(TRAF6 Antibody) at dilution of 1:200



A431 cells were subjected to SDS PAGE followed by western blot with CAF50359(TRAF6 antibody) at dilution of 1:300