

BiomatikTel: (519) 489-7195, (800) 836-8089
Fax: (519) 231-0140, (877) 221-3515
Email: info@ biomatik.com https://www.biomatik.com

anti- HMGCR antibody

Product Information

Catalog No.: CAF50487

Size: 100µg Form: liquid

Purification: Immunogen affinity purified

≥95% as determined by SDS-PAGE Purity:

Host: Rabbit

polyclonal Clonality:

Clone ID: None IsoType: IgG

PBS with 0.02% sodium azide and 50% glycerol pH 7.3, -20°C for 12 Storage:

months (Avoid repeated freeze / thaw cycles.)

Background

HMG-CoA reductase is the rate-limiting enzyme for cholesterol synthesis and is regulated via a negative feedback mechanism mediated by sterols and non-sterol metabolites derived from mevalonate, the product of the reaction catalyzed by reductase. Normally in mammalian cells this enzyme is suppressed by cholesterol derived from the internalization and degradation of low density lipoprotein (LDL) via the LDL receptor. Competitive inhibitors of the reductase induce the expression of LDL receptors in the liver, which in turn increases the catabolism of plasma LDL and lowers the plasma concentration of cholesterol, an important determinant of atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.

Immunogen information

3-hydroxy-3-methylglutaryl-Coenzyme A reductase Immunogen:

HMG CoA reductase, HMGCR, 3-hydroxy-3-methylglutaryl-coenzyme Synonyms:

A reductase, HMDH

Observed MW: 120 kDa Uniprot ID: P04035

Application

Reactivity: Human


Tested Application: ELISA, IHC

Recommended dilution: IHC: 1:20-1:200

Image:



Biomatik Tel: (519) 489-7195, (800) 836-8089 Fax: (519) 231-0140, (877) 221-3515 Email: info@ biomatik.com https://www.biomatik.com



Immunohistochemistry of paraffin-embedded human colon cancer using CAF50487(HMGCR antibody) at dilution of 1:100

