



**CAU24346 Human**  
**Polyclonal Antibody to Cartilage Intermediate Layer Protein (CILP)**  
**Organism Species: *Homo sapiens (Human)***  
***Instruction manual***

FOR RESEARCH USE ONLY  
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

---

12th Edition (Revised in Aug, 2016)



## **[ PROPERTIES ]**

**Source:** Polyclonal antibody preparation

**Host:** Rabbit

**Purification:** Antigen-specific affinity chromatography followed by Protein A affinity chromatography

**Traits:** Liquid

**Concentration:** 0.23mg/ml

**UOM:** 10µg(44µl)

**Applications:** WB; IHC; ICC; IP.

## **[ IMMUNOGEN ]**

**Immunogen:** Recombinant CILP (Ile603~Ala846 (Accession # O75339)) expressed in *E.coli*

**Accession No.:** RPC382Hu01

## **[ APPLICATIONS ]**

Western blotting: 0.5-2µg/mL;1:110-450

Immunohistochemistry: 5-20µg/mL;1:11-45

Immunocytochemistry: 5-20µg/mL;1:11-45

Optimal working dilutions must be determined by end user.

## **[ FORMULATION ]**

**Form & Buffer:** Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300, 50% glycerol.

## **[ STORAGE AND STABILITY ]**

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the



expiration date under appropriate storage condition.

**[ IDENTIFICATION ]**






Figure. Western Blot; Sample:  
Recombinant CILP, Human.



DAB staining on IHC-P; Samples:  
Human Liver Tissue;



DAB staining on IHC-P;  
Samples: Human Liver cancer Tissue

**[ IMPORTANT NOTE ]**

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.