

CAU29275 Human

Monoclonal Antibody to Procollagen III N-Terminal Propeptide (PIIINP)

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



# [PROPERTIES]

Source: Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG2b Kappa

**Purification:** Protein A + Protein G affinity chromatography

Clone number: A6

Traits: Liquid

Concentration: 1mg/ml

**UOM:** 1mg(1000µl)

Applications: WB; IHC; ICC; IP.

### [ IMMUNOGEN ]

Immunogen: Synthetic Peptide, PIIINP conjugated to KLH.

Accession No.: CPA573Hu31

#### [APPLICATIONS]

Western blotting: 0.5-5µg/mL;1:200-2000

Immunohistochemistry: 5-20µg/mL;1:50-200

Immunocytochemistry: 5-20µg/mL;1:50-200

Optimal working dilutions must be determined by end user.

# [FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300,

50% glycerol.

#### [ STORAGE AND STABILITY ]

**Storage:** Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.

Aliquot and store at -20°C for 12 months.

**Stability Test:** The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no



obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

# [ IDENTIFICATION ]





Figure. DAB staining on IHC-P; Samples: Human Liver Tissue.



Western Blot; Sample: Human Hela cell lysate;




Figure. DAB staining on IHC-P; Samples: Human Kidney Tissue.

Primary Ab: 5µg/ml Mouse Anti-Human
PIIINP Antibody

Second Ab: 0.2µg/mL HRP-Linked
Caprine Anti-Mouse IgG Polyclonal
Antibody

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.